

Acta Crystallographica Section E

Structure Reports
Online

ISSN 1600-5368

Editors: W. T. A. Harrison, H. Stoeckli-Evans,
E. R. T. Tiekink and M. Weil

Methyl (*E*)-2-[(3-chloro-4-cyanophenyl)imino]-4-(4-chlorophenyl)-6-methyl-1,2,3,4-tetrahydropyrimidine-5-carboxylate

K. N. Venugopala, Susanta K. Nayak and Bharti Odhav

Acta Cryst. (2012). E68, o2977–o2978

This open-access article is distributed under the terms of the Creative Commons Attribution Licence <http://creativecommons.org/licenses/by/2.0/uk/legalcode>, which permits unrestricted use, distribution, and reproduction in any medium, provided the original authors and source are cited.



Acta Crystallographica Section E: Structure Reports Online is the IUCr's highly popular open-access structural journal. It provides a simple and easily accessible publication mechanism for the growing number of inorganic, metal-organic and organic crystal structure determinations. The electronic submission, validation, refereeing and publication facilities of the journal ensure very rapid and high-quality publication, whilst key indicators and validation reports provide measures of structural reliability. The journal publishes over 4000 structures per year. The average publication time is less than one month.

Crystallography Journals Online is available from journals.iucr.org

Methyl (*E*)-2-[(3-chloro-4-cyanophenyl)-imino]-4-(4-chlorophenyl)-6-methyl-1,2,3,4-tetrahydropyrimidine-5-carboxylate

K. N. Venugopala,^{a*} Susanta K. Nayak^{b*} and Bharti Odhav^a

^aDepartment of Biotechnology and Food Technology, Durban University of Technology, Durban 4001, South Africa, and ^bCenter for Nano Science and Technology @ Polimi, Istituto Italiano di Tecnologia, Via Pascoli 70/3, 20133 Milan, Italy

Correspondence e-mail: kharigattav@dut.ac.za, nksusa@gmail.com


Received 5 September 2012; accepted 16 September 2012

Key indicators: single-crystal X-ray study; $T = 173$ K; mean $\sigma(\text{C}-\text{C}) = 0.004$ Å; disorder in main residue; R factor = 0.046; wR factor = 0.130; data-to-parameter ratio = 13.5.

In the title compound, $\text{C}_{20}\text{H}_{16}\text{Cl}_2\text{N}_4\text{O}_2$, the dihedral angles between the planes of the chlorophenyl, chlorocyanophenyl-imine and ester groups and the plane of the six-membered tetrahydropyrimidine ring are 86.9 (2), 72.6 (2) and 7.9 (2)°, respectively. The Cl atom substituent on the cyanophenyl ring is disordered over two rotationally related sites [occupancy factors 0.887 (2):0.113 (2)], while the molecular conformation is stabilized by the presence of an intramolecular aromatic C–H... π interaction. Both N–H groups participate in separate intermolecular hydrogen-bonding associations with centrosymmetric cyclic motifs [graph sets $R_2^2(8)$ and $R_2^2(12)$], resulting in ribbons parallel to [010]. Further weak C–H...O hydrogen bonds link these ribbons into a two-dimensional molecular assembly.

Related literature

For crystal structures of the dihydropyrimidines, see: Nayak *et al.* (2010); Nayak, Venugopala, Govender *et al.* (2011); Nayak, Venugopala, Chopra & Guru Row (2011). For background on the applications of dihydropyrimidines, see: Kappe (2000). For graph-set analysis, see: Bernstein *et al.* (1995).



Experimental

Crystal data

$\text{C}_{20}\text{H}_{16}\text{Cl}_2\text{N}_4\text{O}_2$

$M_r = 415.27$

Monoclinic, $P2_1/c$

$a = 11.905$ (8) Å

$b = 13.729$ (9) Å

$c = 12.782$ (8) Å

$\beta = 108.366$ (14)°

$V = 1983$ (2) Å³

$Z = 4$

Mo $K\alpha$ radiation

$\mu = 0.35$ mm⁻¹

$T = 173$ K

$0.23 \times 0.12 \times 0.03$ mm

Data collection

Bruker Kappa DUO APEXII

diffractometer

Absorption correction: multi-scan

(*SADABS*; Bruker, 2008)

$T_{\min} = 0.924$, $T_{\max} = 0.990$

9454 measured reflections

3497 independent reflections

2324 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.029$

Standard reflections: 0

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.046$

$wR(F^2) = 0.130$

$S = 1.01$

3497 reflections

259 parameters

H-atom parameters constrained

$\Delta\rho_{\text{max}} = 0.41$ e Å⁻³

$\Delta\rho_{\text{min}} = -0.37$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

Cg1 is the mid-point of the C3=C4 bond.

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
N1–H1 \cdots N4A'	0.88	2.21	2.981 (4)	147
N2–H2 \cdots N3 ⁱⁱ	0.88	2.09	2.966 (4)	172
C15A–H15A \cdots O1 ⁱⁱⁱ	0.95	2.39	3.322 (4)	169
C12–H12 \cdots Cg1	0.95	2.85	3.290 (2)	109

Symmetry codes: (i) $-x + 2, -y, -z$; (ii) $-x + 2, -y + 1, -z$; (iii) $-x + 2, y, -z + \frac{1}{2}$.

Data collection: *APEX2* (Bruker, 2008); cell refinement: *SAINT* (Bruker, 2008); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3 for Windows* (Farrugia, 1997) and *Mercury* (Macrae *et al.*, 2008); software used to prepare material for publication: *PLATON* (Spek, 2009) and *PARST* (Nardelli, 1995).

The authors thank Durban University of Technology for facilities. KNV thanks NRF South Africa for a DST/NRF Innovation Postdoctoral Fellowship.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: ZS2233).

References

- Bernstein, J., Davis, R. E., Shimoni, L. & Chang, N.-L. (1995). *Angew. Chem. Int. Ed. Engl.* **34**, 1555–1573.
- Bruker (2008). *APEX2, SAINT and SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
- Kappe, C. O. (2000). *Eur. J. Med. Chem.* **35**, 1043–1052.
- Macrae, C. F., Bruno, I. J., Chisholm, J. A., Edgington, P. R., McCabe, P., Pidcock, E., Rodriguez-Monge, L., Taylor, R., van de Streek, J. & Wood, P. A. (2008). *J. Appl. Cryst.* **41**, 466–470.
- Nardelli, M. (1995). *J. Appl. Cryst.* **28**, 659.
- Nayak, S. K., Venugopala, K. N., Chopra, D. & Guru Row, T. N. (2011). *CrystEngComm*, **13**, 591–605.
- Nayak, S. K., Venugopala, K. N., Chopra, D., Vasu & Guru Row, T. N. (2010). *CrystEngComm*, **12**, 1205–1216.
- Nayak, S. K., Venugopala, K. N., Govender, T., Kruger, H. G., Maguire, G. E. M. & Row, T. N. G. (2011). *Acta Cryst. E* **67**, o3069–o3070.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Spek, A. L. (2009). *Acta Cryst. D* **65**, 148–155.

supplementary materials

Acta Cryst. (2012). E68, o2977–o2978 [doi:10.1107/S1600536812039451]

Methyl (*E*)-2-[(3-chloro-4-cyanophenyl)imino]-4-(4-chlorophenyl)-6-methyl-1,2,3,4-tetrahydropyrimidine-5-carboxylate

K. N. Venugopala, Susanta K. Nayak and Bharti Odhav

Comment

The multifunctionalized dihydropyrimidones (DHPMs) are prime target molecules for their therapeutic and pharmacological properties (Kappe, 2000). Due to the vast range of applications of this class of compounds we have been investigating conformational and packing features of tetrahydropyrimidine derivatives of this title compound (Nayak *et al.*, 2010; Nayak, Venugopala, Govender *et al.*, 2011; Nayak, Venugopala, Chopra & Guru Row (2011). In a continuation of our work on synthesis of heterocyclic compounds for biological properties, herein we report the single-crystal structure of the title compound, C₂₀H₁₆Cl₂N₄O₂.

In this molecule (Fig. 1), the dihedral angles between the planes of the 4-chlorophenyl, 3-chloro-4-cyanophenylimino and ester groups (O2/C2/O1/C1) and the plane of the six-membered tetrahydropyrimidine ring are 86.9 (2)°, 72.6 (2)° and 7.9 (2)° respectively. The conformation of the molecule is stabilized by an intra-molecular C—H... π interaction (Table 1) wherein the aryl hydrogen H12 is oriented towards the π electrons of the C3=C4 bond. The *meta*-related chlorine substituent on the cyanophenyl ring is disordered over two rotationally-related sites [occupancy factors 0.887 (2) (*A*): 0.113 (2) (*B*)]. Both N—H groups participate in separate intermolecular hydrogen-bonding associations giving centrosymmetric cyclic motifs [graph sets $R_2^2(8)$ and $R_2^2(12)$ (Bernstein *et al.*, 1995)], resulting in ribbons parallel to [010] (Fig. 2*a*). Further weak C—H...O hydrogen bonds (Fig. 2*b*) link these ribbons into a two-dimensional molecular assembly. Present also is a short intermolecular Cl...Cl interaction [C11...Cl2Bⁱⁱⁱ; 2.884 (7) Å (symmetry code $-x + 1, y, -z - 1/2$)].

Experimental

A mixture of methyl-2-chloro-4-(*p*-chlorophenyl)-6-methyl-1,4-dihydropyrimidine-5-carboxylate (1 mmol), 4-amino-2-chlorobenzonitrile (1 mmol) and methanamine (1 mmol) in 2-propanol (5 ml) was refluxed for 10 h. The reaction completion was monitored by TLC. The reaction medium was cooled to room temperature, the product was filtered, washed with cold 2-propanol and dried to obtain the crude product. The product was purified by recrystallization using ethanol in 69% yield as a yellow solid (m.p. 431 (2) K). Crystals suitable for single-crystal X-ray study were obtained from methanol solvent using slow evaporation at room temperature.

Refinement

The 3-chloro-4-cyanophenylimino group was treated as disordered over two possible rotation-related sites (*A* and *B*), having refined site occupancy factors of 0.887 (2) and 0.113 (2), respectively. All H atoms were positioned geometrically with N—H = 0.88 Å, C—H = 0.95–1.00 Å and refined using a riding model with $U_{iso}(H) = 1.2U_{eq}(C/N)$ except for the methyl group where $U_{iso}(H) = 1.5U_{eq}(C)$.

Computing details

Data collection: *APEX2* (Bruker, 2008); cell refinement: *SAINT* (Bruker, 2008); data reduction: *SAINT* (Bruker, 2008); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3 for Windows* (Farrugia, 1997) and *Mercury* (Macrae *et al.*, 2008); software used to prepare material for publication: *PLATON* (Spek, 2009) and *PARST* (Nardelli, 1995).





Figure 1

A view of the title compound with the atom numbering scheme and displacement ellipsoids for non-H atoms drawn at the 50% probability level. The intramolecular C—H... π interaction is shown as dashed lines. The disordered chlorine positions are differentiated as *A* and *B*.


Figure 2

(a) Intermolecular N—H...N hydrogen-bonding associations form an infinite ribbon structure. (b) Further C—H...O hydrogen bonds link the ribbons giving a two-dimensional network structure.

Methyl (*E*)-2-[(3-chloro-4-cyanophenyl)imino]-4-(4-chlorophenyl)-6-methyl-1,2,3,4-tetrahydropyrimidine-5-carboxylate

Crystal data

$C_{20}H_{16}Cl_2N_4O_2$

$M_r = 415.27$

Monoclinic, $P2_1/c$

Hall symbol: $-P\ 2_1/c$

$a = 11.905\ (8)\ \text{\AA}$

$b = 13.729\ (9)\ \text{\AA}$

$c = 12.782\ (8)\ \text{\AA}$

$\beta = 108.366\ (14)^\circ$

$V = 1983\ (2)\ \text{\AA}^3$

$Z = 4$

$F(000) = 856$

$D_x = 1.391\ \text{Mg m}^{-3}$

Melting point: 431(2) K

Mo $K\alpha$ radiation, $\lambda = 0.71073\ \text{\AA}$

Cell parameters from 650 reflections

$\theta = 1.5\text{--}25.0^\circ$

$\mu = 0.35\ \text{mm}^{-1}$

$T = 173\ \text{K}$

Plate, yellow

$0.23 \times 0.12 \times 0.03\ \text{mm}$

Data collection

Bruker Kappa DUO APEXII

diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

$0.5^\circ\ \varphi$ scans and ω scans

Absorption correction: multi-scan

(*SADABS*; Bruker, 2008)

$T_{\min} = 0.924$, $T_{\max} = 0.990$

9454 measured reflections

3497 independent reflections

2324 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.029$

$\theta_{\max} = 25.0^\circ$, $\theta_{\min} = 1.5^\circ$

$h = -14 \rightarrow 13$

$k = -16 \rightarrow 16$

$l = -7 \rightarrow 15$

Refinement

Refinement on F^2
 Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.046$
 $wR(F^2) = 0.130$
 $S = 1.01$
 3497 reflections
 259 parameters
 0 restraints
 Primary atom site location: structure-invariant
 direct methods

Secondary atom site location: difference Fourier
 map
 Hydrogen site location: inferred from
 neighbouring sites
 H-atom parameters constrained
 $w = 1/[\sigma^2(F_o^2) + (0.0628P)^2 + 0.7735P]$
 where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} = 0.001$
 $\Delta\rho_{\max} = 0.41 \text{ e } \text{\AA}^{-3}$
 $\Delta\rho_{\min} = -0.37 \text{ e } \text{\AA}^{-3}$

Special details

Geometry. All s.u.'s (except the s.u. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell s.u.'s are taken into account individually in the estimation of s.u.'s in distances, angles and torsion angles; correlations between s.u.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell s.u.'s is used for estimating s.u.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > 2\sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$	Occ. (<1)
Cl2A	1.27478 (8)	0.07230 (7)	0.07772 (9)	0.0732 (3)	0.8872 (16)
C14A	1.0144 (2)	0.26844 (18)	-0.0167 (2)	0.0429 (7)	0.8872 (16)
C15A	1.1235 (3)	0.2245 (2)	0.0335 (2)	0.0511 (7)	0.8872 (16)
H15A	1.1837	0.2590	0.0874	0.061*	0.8872 (16)
C16A	1.1443 (2)	0.1307 (2)	0.0047 (2)	0.0490 (7)	0.8872 (16)
C17A	1.0592 (2)	0.07963 (17)	-0.0761 (2)	0.0387 (6)	0.8872 (16)
C18A	0.9496 (2)	0.12388 (18)	-0.1256 (2)	0.0424 (6)	0.8872 (16)
H18A	0.8900	0.0898	-0.1806	0.051*	0.8872 (16)
C19A	0.9270 (2)	0.21625 (19)	-0.0957 (2)	0.0435 (6)	0.8872 (16)
H19A	0.8514	0.2448	-0.1291	0.052*	0.8872 (16)
C20A	1.0824 (3)	-0.0174 (2)	-0.1067 (2)	0.0449 (7)	0.8872 (16)
N4A	1.0998 (2)	-0.09457 (18)	-0.1320 (2)	0.0584 (7)	0.8872 (16)
Cl2B	0.8417 (6)	0.0473 (6)	-0.1826 (7)	0.0732 (3)	0.1128 (16)
C14B	1.0144 (2)	0.26844 (18)	-0.0167 (2)	0.0429 (7)	0.1128 (16)
C15B	1.1235 (3)	0.2245 (2)	0.0335 (2)	0.0511 (7)	0.1128 (16)
H15B	1.1837	0.2590	0.0874	0.061*	0.1128 (16)
C16B	1.1443 (2)	0.1307 (2)	0.0047 (2)	0.0490 (7)	0.1128 (16)
H16B	1.2182	0.1007	0.0410	0.059*	0.1128 (16)
C17B	1.0592 (2)	0.07963 (17)	-0.0761 (2)	0.0387 (6)	0.1128 (16)
C18B	0.9496 (2)	0.12388 (18)	-0.1256 (2)	0.0424 (6)	0.1128 (16)
C19B	0.9270 (2)	0.21625 (19)	-0.0957 (2)	0.0435 (6)	0.1128 (16)
H19B	0.8514	0.2448	-0.1291	0.052*	0.1128 (16)
C20B	1.0824 (3)	-0.0174 (2)	-0.1067 (2)	0.0449 (7)	0.1128 (16)
N4B	1.0998 (2)	-0.09457 (18)	-0.1320 (2)	0.0584 (7)	0.1128 (16)
Cl1	0.35886 (9)	0.15552 (9)	-0.18125 (10)	0.1015 (4)	
O1	0.67082 (15)	0.36934 (13)	0.30483 (14)	0.0440 (5)	

O2	0.6947 (2)	0.53124 (14)	0.30330 (18)	0.0635 (6)
N1	0.88569 (18)	0.31127 (15)	0.12045 (18)	0.0418 (5)
H1	0.9209	0.2541	0.1288	0.050*
N2	0.89937 (19)	0.47517 (14)	0.08817 (18)	0.0445 (6)
H2	0.9239	0.5226	0.0544	0.053*
N3	0.9962 (2)	0.36677 (15)	0.0096 (2)	0.0480 (6)
C1	0.6064 (3)	0.3833 (2)	0.3825 (2)	0.0545 (8)
H1A	0.6603	0.4078	0.4523	0.082*
H1B	0.5725	0.3211	0.3952	0.082*
H1C	0.5426	0.4305	0.3525	0.082*
C2	0.7139 (2)	0.4517 (2)	0.2724 (2)	0.0429 (6)
C3	0.7800 (2)	0.42850 (18)	0.1967 (2)	0.0393 (6)
C4	0.8352 (2)	0.49915 (19)	0.1574 (2)	0.0405 (6)
C5	0.8350 (3)	0.60622 (18)	0.1828 (2)	0.0478 (7)
H5A	0.7538	0.6311	0.1557	0.072*
H5B	0.8843	0.6411	0.1466	0.072*
H5C	0.8667	0.6161	0.2627	0.072*
C6	0.7861 (2)	0.32283 (18)	0.1639 (2)	0.0398 (6)
H6	0.8027	0.2821	0.2320	0.048*
C7	0.6733 (2)	0.28406 (18)	0.0795 (2)	0.0403 (6)
C8	0.6315 (3)	0.1926 (2)	0.0943 (3)	0.0552 (8)
H8	0.6691	0.1570	0.1597	0.066*
C9	0.5350 (3)	0.1527 (2)	0.0139 (3)	0.0697 (10)
H9	0.5076	0.0894	0.0234	0.084*
C10	0.4795 (3)	0.2057 (3)	-0.0795 (3)	0.0625 (9)
C11	0.5174 (3)	0.2981 (2)	-0.0955 (3)	0.0589 (8)
H11	0.4778	0.3345	-0.1598	0.071*
C12	0.6147 (3)	0.3362 (2)	-0.0151 (2)	0.0496 (7)
H12	0.6420	0.3994	-0.0250	0.060*
C13	0.9271 (2)	0.38082 (17)	0.0692 (2)	0.0400 (6)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C12A	0.0432 (5)	0.0708 (6)	0.0957 (7)	0.0139 (4)	0.0076 (5)	-0.0130 (5)
C14A	0.0484 (16)	0.0313 (14)	0.0634 (18)	-0.0036 (13)	0.0384 (15)	-0.0031 (13)
C15A	0.0442 (16)	0.0458 (16)	0.0676 (19)	-0.0057 (14)	0.0237 (15)	-0.0131 (14)
C16A	0.0393 (15)	0.0455 (16)	0.0652 (18)	0.0060 (13)	0.0207 (14)	-0.0046 (14)
C17A	0.0473 (16)	0.0295 (13)	0.0491 (15)	0.0024 (12)	0.0290 (13)	-0.0010 (12)
C18A	0.0447 (16)	0.0343 (14)	0.0514 (16)	-0.0026 (12)	0.0197 (13)	-0.0027 (12)
C19A	0.0426 (15)	0.0337 (14)	0.0596 (17)	0.0016 (13)	0.0238 (14)	0.0004 (13)
C20A	0.0529 (17)	0.0394 (16)	0.0462 (16)	0.0063 (13)	0.0214 (13)	0.0014 (13)
N4A	0.0752 (18)	0.0419 (15)	0.0620 (16)	0.0152 (13)	0.0273 (14)	-0.0020 (12)
C12B	0.0432 (5)	0.0708 (6)	0.0957 (7)	0.0139 (4)	0.0076 (5)	-0.0130 (5)
C14B	0.0484 (16)	0.0313 (14)	0.0634 (18)	-0.0036 (13)	0.0384 (15)	-0.0031 (13)
C15B	0.0442 (16)	0.0458 (16)	0.0676 (19)	-0.0057 (14)	0.0237 (15)	-0.0131 (14)
C16B	0.0393 (15)	0.0455 (16)	0.0652 (18)	0.0060 (13)	0.0207 (14)	-0.0046 (14)
C17B	0.0473 (16)	0.0295 (13)	0.0491 (15)	0.0024 (12)	0.0290 (13)	-0.0010 (12)
C18B	0.0447 (16)	0.0343 (14)	0.0514 (16)	-0.0026 (12)	0.0197 (13)	-0.0027 (12)
C19B	0.0426 (15)	0.0337 (14)	0.0596 (17)	0.0016 (13)	0.0238 (14)	0.0004 (13)

C20B	0.0529 (17)	0.0394 (16)	0.0462 (16)	0.0063 (13)	0.0214 (13)	0.0014 (13)
N4B	0.0752 (18)	0.0419 (15)	0.0620 (16)	0.0152 (13)	0.0273 (14)	-0.0020 (12)
Cl1	0.0658 (6)	0.1042 (8)	0.1193 (8)	-0.0215 (6)	0.0072 (6)	-0.0381 (7)
O1	0.0411 (10)	0.0440 (11)	0.0553 (11)	0.0003 (9)	0.0270 (9)	0.0015 (9)
O2	0.0846 (16)	0.0439 (12)	0.0850 (15)	0.0025 (11)	0.0598 (14)	-0.0096 (11)
N1	0.0393 (12)	0.0288 (11)	0.0661 (14)	0.0043 (9)	0.0293 (11)	-0.0007 (10)
N2	0.0528 (14)	0.0280 (11)	0.0687 (15)	-0.0037 (10)	0.0420 (13)	-0.0065 (10)
N3	0.0545 (14)	0.0296 (11)	0.0752 (16)	-0.0041 (11)	0.0423 (13)	-0.0089 (11)
C1	0.0521 (17)	0.064 (2)	0.0577 (18)	-0.0015 (15)	0.0325 (15)	0.0001 (15)
C2	0.0380 (14)	0.0409 (15)	0.0544 (16)	0.0038 (12)	0.0213 (13)	-0.0015 (13)
C3	0.0371 (14)	0.0342 (14)	0.0518 (16)	0.0005 (12)	0.0214 (12)	-0.0028 (12)
C4	0.0381 (14)	0.0353 (14)	0.0538 (16)	0.0002 (12)	0.0225 (13)	-0.0072 (12)
C5	0.0488 (16)	0.0350 (14)	0.0701 (19)	-0.0037 (13)	0.0339 (15)	-0.0100 (13)
C6	0.0392 (14)	0.0331 (13)	0.0563 (16)	0.0023 (11)	0.0283 (13)	0.0018 (12)
C7	0.0374 (14)	0.0331 (14)	0.0599 (17)	0.0009 (12)	0.0287 (13)	-0.0054 (13)
C8	0.0443 (16)	0.0344 (15)	0.089 (2)	0.0015 (13)	0.0237 (16)	0.0053 (15)
C9	0.0474 (18)	0.0366 (17)	0.123 (3)	-0.0069 (15)	0.024 (2)	-0.0123 (19)
C10	0.0458 (18)	0.061 (2)	0.081 (2)	-0.0060 (17)	0.0200 (17)	-0.0222 (19)
C11	0.0559 (19)	0.071 (2)	0.0545 (18)	-0.0044 (17)	0.0248 (16)	-0.0050 (16)
C12	0.0520 (17)	0.0454 (16)	0.0595 (18)	-0.0070 (14)	0.0290 (15)	-0.0006 (14)
C13	0.0384 (14)	0.0288 (13)	0.0592 (16)	-0.0039 (11)	0.0246 (13)	-0.0074 (12)

Geometric parameters (Å, °)

Cl2A—C16A	1.736 (3)	N3—C13	1.300 (3)
C14A—C15A	1.392 (4)	C1—H1A	0.9800
C14A—C19A	1.398 (4)	C1—H1B	0.9800
C14A—N3	1.424 (3)	C1—H1C	0.9800
C15A—C16A	1.382 (4)	C2—C3	1.461 (3)
C15A—H15A	0.9500	C3—C4	1.355 (3)
C16A—C17A	1.388 (4)	C3—C6	1.518 (4)
C17A—C18A	1.397 (4)	C4—C5	1.506 (4)
C17A—C20A	1.439 (4)	C5—H5A	0.9800
C18A—C19A	1.375 (4)	C5—H5B	0.9800
C18A—H18A	0.9500	C5—H5C	0.9800
C19A—H19A	0.9500	C6—C7	1.530 (4)
C20A—N4A	1.146 (3)	C6—H6	1.0000
Cl1—C10	1.747 (3)	C7—C8	1.385 (4)
O1—C2	1.359 (3)	C7—C12	1.389 (4)
O1—C1	1.446 (3)	C8—C9	1.389 (4)
O2—C2	1.207 (3)	C8—H8	0.9500
N1—C13	1.337 (3)	C9—C10	1.376 (5)
N1—C6	1.468 (3)	C9—H9	0.9500
N1—H1	0.8800	C10—C11	1.383 (5)
N2—C13	1.377 (3)	C11—C12	1.386 (4)
N2—C4	1.378 (3)	C11—H11	0.9500
N2—H2	0.8800	C12—H12	0.9500
C15A—C14A—C19A	119.0 (2)	C2—C3—C6	118.3 (2)
C15A—C14A—N3	119.4 (3)	C3—C4—N2	120.0 (2)

C19A—C14A—N3	121.5 (3)	C3—C4—C5	125.8 (2)
C16A—C15A—C14A	120.0 (3)	N2—C4—C5	114.3 (2)
C16A—C15A—H15A	120.0	C4—C5—H5A	109.5
C14A—C15A—H15A	120.0	C4—C5—H5B	109.5
C15A—C16A—C17A	121.3 (3)	H5A—C5—H5B	109.5
C15A—C16A—C12A	119.5 (2)	C4—C5—H5C	109.5
C17A—C16A—C12A	119.1 (2)	H5A—C5—H5C	109.5
C16A—C17A—C18A	118.4 (2)	H5B—C5—H5C	109.5
C16A—C17A—C20A	120.9 (3)	N1—C6—C3	108.84 (19)
C18A—C17A—C20A	120.7 (2)	N1—C6—C7	109.2 (2)
C19A—C18A—C17A	120.8 (3)	C3—C6—C7	114.8 (2)
C19A—C18A—H18A	119.6	N1—C6—H6	107.9
C17A—C18A—H18A	119.6	C3—C6—H6	107.9
C18A—C19A—C14A	120.5 (3)	C7—C6—H6	107.9
C18A—C19A—H19A	119.8	C8—C7—C12	118.8 (3)
C14A—C19A—H19A	119.8	C8—C7—C6	119.5 (3)
N4A—C20A—C17A	179.3 (4)	C12—C7—C6	121.6 (2)
C2—O1—C1	115.6 (2)	C7—C8—C9	120.3 (3)
C13—N1—C6	125.1 (2)	C7—C8—H8	119.8
C13—N1—H1	117.5	C9—C8—H8	119.8
C6—N1—H1	117.5	C10—C9—C8	119.5 (3)
C13—N2—C4	123.3 (2)	C10—C9—H9	120.3
C13—N2—H2	118.3	C8—C9—H9	120.3
C4—N2—H2	118.3	C9—C10—C11	121.6 (3)
C13—N3—C14A	116.7 (2)	C9—C10—C11	119.7 (3)
O1—C1—H1A	109.5	C11—C10—C11	118.8 (3)
O1—C1—H1B	109.5	C10—C11—C12	118.1 (3)
H1A—C1—H1B	109.5	C10—C11—H11	120.9
O1—C1—H1C	109.5	C12—C11—H11	120.9
H1A—C1—H1C	109.5	C11—C12—C7	121.6 (3)
H1B—C1—H1C	109.5	C11—C12—H12	119.2
O2—C2—O1	121.6 (2)	C7—C12—H12	119.2
O2—C2—C3	127.6 (2)	N3—C13—N1	125.5 (2)
O1—C2—C3	110.7 (2)	N3—C13—N2	118.3 (2)
C4—C3—C2	121.0 (2)	N1—C13—N2	116.1 (2)
C4—C3—C6	120.7 (2)		
C19A—C14A—C15A—C16A	0.1 (4)	C13—N1—C6—C3	-29.3 (3)
N3—C14A—C15A—C16A	-176.8 (2)	C13—N1—C6—C7	96.7 (3)
C14A—C15A—C16A—C17A	1.9 (4)	C4—C3—C6—N1	18.4 (3)
C14A—C15A—C16A—C12A	-173.4 (2)	C2—C3—C6—N1	-161.1 (2)
C15A—C16A—C17A—C18A	-2.3 (4)	C4—C3—C6—C7	-104.4 (3)
C12A—C16A—C17A—C18A	173.05 (19)	C2—C3—C6—C7	76.1 (3)
C15A—C16A—C17A—C20A	179.0 (2)	N1—C6—C7—C8	101.1 (3)
C12A—C16A—C17A—C20A	-5.7 (4)	C3—C6—C7—C8	-136.3 (2)
C16A—C17A—C18A—C19A	0.6 (4)	N1—C6—C7—C12	-75.4 (3)
C20A—C17A—C18A—C19A	179.4 (2)	C3—C6—C7—C12	47.1 (3)
C17A—C18A—C19A—C14A	1.3 (4)	C12—C7—C8—C9	2.1 (4)
C15A—C14A—C19A—C18A	-1.7 (4)	C6—C7—C8—C9	-174.6 (2)

N3—C14A—C19A—C18A	175.2 (2)	C7—C8—C9—C10	-1.5 (5)
C15A—C14A—N3—C13	-108.4 (3)	C8—C9—C10—C11	0.0 (5)
C19A—C14A—N3—C13	74.8 (3)	C8—C9—C10—C11	179.5 (2)
C1—O1—C2—O2	-3.0 (4)	C9—C10—C11—C12	0.8 (5)
C1—O1—C2—C3	178.4 (2)	C11—C10—C11—C12	-178.7 (2)
O2—C2—C3—C4	4.3 (5)	C10—C11—C12—C7	-0.2 (4)
O1—C2—C3—C4	-177.2 (2)	C8—C7—C12—C11	-1.2 (4)
O2—C2—C3—C6	-176.2 (3)	C6—C7—C12—C11	175.3 (2)
O1—C2—C3—C6	2.3 (3)	C14A—N3—C13—N1	10.1 (4)
C2—C3—C4—N2	178.5 (2)	C14A—N3—C13—N2	-174.5 (2)
C6—C3—C4—N2	-1.0 (4)	C6—N1—C13—N3	-163.9 (3)
C2—C3—C4—C5	-1.0 (4)	C6—N1—C13—N2	20.6 (4)
C6—C3—C4—C5	179.5 (3)	C4—N2—C13—N3	-174.4 (3)
C13—N2—C4—C3	-10.6 (4)	C4—N2—C13—N1	1.4 (4)
C13—N2—C4—C5	168.9 (3)		

Hydrogen-bond geometry (\AA , $^\circ$)

Cg1 is the midpoint of the C3=C4 bond. [Please check added text]

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
N1—H1 \cdots N4A ⁱ	0.88	2.21	2.981 (4)	147
N2—H2 \cdots N3 ⁱⁱ	0.88	2.09	2.966 (4)	172
C15A—H15A \cdots O1 ⁱⁱⁱ	0.95	2.39	3.322 (4)	169
C12—H12 \cdots Cg1	0.95	2.85	3.290 (2)	109

Symmetry codes: (i) $-x+2, -y, -z$; (ii) $-x+2, -y+1, -z$; (iii) $-x+2, y, -z+1/2$.